



Empirical Mode Decomposition and its application to Palm Tree Mode analysis

Maszl, Ch.; Hoffmann, Mathias; Naulin, Volker; Juul Rasmussen, Jens; Gennrich, F.; Schrittwieser, R.

Published in:
Proceedings

Publication date:
2011

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Maszl, C., Hoffmann, M., Naulin, V., Juul Rasmussen, J., Gennrich, F., & Schrittwieser, R. (2011). Empirical Mode Decomposition and its application to Palm Tree Mode analysis. In *Proceedings* (pp. P4.064)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Empirical Mode Decomposition and its application to Palm Tree Mode analysis

Ch. Maszl¹, M. Hoffmann², V. Naulin², J. J. Rasmussen², F. Gennrich¹,

R. Schrittwieser¹ and JET EFDA Contributors*

JET-EFDA, Culham Science Centre, OX14 3DB, Abingdon, UK

¹*Association EURATOM/ÖAW, Institute for Ion Physics and Applied Physics,*

University of Innsbruck, Austria

²*Association EURATOM/RISØ DTU, Technical University of Denmark, Roskilde, Denmark*

Introduction

Recent works report the existence of coherent long-lived structures or current filaments in the edge of tokamak plasmas. These are found as confined current ribbons (outer modes) [1] or palm tree modes (PTM) [2] in the edge of JET for example. An intuitive model was proposed in [3] which describes the genesis of these structures by a blob/hole creation mechanism during the ELM. ELM filaments with excess current [4], temperature and density are treated as blobs whereas it is believed that they leave corresponding holes behind due to current conservation considerations on short time scales [5]. It therefore complements the MHD description by a blob/hole or "quasi-particle" picture.

From pellet injection experiments in DIII-D and Tore Supra it is known that maxima of the mass deposition profiles are close to the magnetic $q = 2$ and $q = 3$ surfaces. Low order rational surfaces appear to slow down or even stop the polarization drift towards the low field side of the plasma [6]. Similarly




Figure 1: Illustration of an ELM in the exhaust phase at the outboard side of a tokamak. If an ELM filament (white) is stopped and captured at e.g. the $q = 4$ surface (green) an outer mode or current ribbon could be created [1]. Similarly a hole (black) captured on a $q = 3$ surface could create a palm tree mode (PTM) [2]. PTMs and outer modes are possibly "twins".

*see the Appendix of F. Romanelli et al., Proceedings of the 23rd IAEA Fusion Energy Conference 2010, Daejeon, Korea

it is to be expected that the lifetime of blobs and holes is increased in the vicinity of rational surfaces. The ∇B polarisation of holes is opposite to blobs. Under the assumption that the interchange drive of a polarised ELM-induced hole moves it into the proximity of a rational surface with low q , it should be slowed down or even stopped. Furthermore, large holes and blobs could close on themselves in the vicinity of a rational magnetic field line. For a closed filament, charge interchange imbalance would be short-circuited by parallel currents which would stop the filament completely. The result would be a closed, well localized filament with a significantly increased lifetime which can only be filled by slow perpendicular transport. Fig. 1 illustrates the proposed events before the holes and blobs reach the resonant surfaces. In this way of thinking outer modes are ELM filaments which were not able to propagate to the scrape-off layer whereas palm tree modes are signatures of ELM-induced holes. Nevertheless, the probability to create PTMs is smaller because holes travel up the temperature gradient into regions with increasing parallel transport. If this is true one can ask why the spectra of outer modes and PTMs show rich harmonics in the Fourier spectra? Fourier spectral analysis uses linear superpositions of unlocalised trigonometric functions. Therefore FFT can only be applied to linear and stationary time series. If nonlinear effects deform the wave profiles towards localisation, FFT needs additional harmonic components to describe these profiles. Non-stationarity and nonlinearity can induce spurious harmonic components that cause energy spreading. An example for such a behaviour would be a Fourier decomposition of a flash light, represented by a delta function in the time domain. Localised in time functions produce less localised FFTs. Spatial localised structures are therefore not independently traveling waves, but highly coherent wave packages. The non-stationary time series may be represented as follows [7].

$$X(t) = \sum_{j=1}^n a_j(t) \exp(i \int \omega_j(t) dt) \quad (1)$$

In order to extract signals from nonlinear or non-stationary time series $X(t)$, the adaptive time-frequency data analysis method "Empirical Mode Decomposition" (EMD) [7] is suitable. EMD acts as a dyadic filter bank to decompose the series into a finite set of so-called Intrinsic Mode Functions (IMF). Each IMF reflects the data on a different time-scale and admits well-behaved Hilbert transforms. EMD enables examinations of noisy linear and nonlinear processes. In contrast to Fourier spectral analysis the amplitudes $a_j(t)$ and frequencies $\omega_j(t)$ obtained by EMD are functions of time (Eq. 1). Amplitude and frequency modulations described by the IMFs are now clearly separated.

Results and Discussion

Phenomena like PTMs exhibit a current, density and a temperature perturbation. Therefore EMD was applied to MHD and ECE data of JET. Electron temperature fluctuations of the PTM measured by ECE show different behaviour in different channels of the diagnostic (Fig. 2 a). The investigated channels B2:001, B1:012 and B1:011 cover parts of the pedestal region of JET for pulse #52011, $t = 19.164 - 19.166$ s. Channel B1:011 (green) shows large positive temperature spikes as the PTM filament enters the detection volume. In contrast, channel B2:001 (orange) shows negative spikes compared to the background temperature. Channel B1:012 (blue) suggests a symmetric modulation of the electron temperature without temperature spikes as in channel B1:011 or B2:001. This observation was quantified in PDFs.




Figure 2: a) Electron temperature fluctuations measured by three different ECE channels at different radial positions. The vertical red line gives insight into the phase relation between the three signals (JET Pulse: #52011, $t = 19.164 - 19.166$ s). b) EMD of channel B1:011. The red signal shows spiky temperature peaks slightly advanced with respect to the maxima of the black line. The black line gives the fraction of the current perturbation. A superposition of these two signals yields the detrended green measured signal from a).

Fig. 2 shows the result of EMD. The red signal is the superposition of the first five IMFs, where the black line shows the detrended raw data (green, Fig. 2) minus the red signals. Although ECE data sets are pretty noisy, signs of temperatures spikes can be observed in the red signal. It is assumed that these spikes represent the temperature inside the filament. The black line is interpreted as distortion of the local equilibrium due to the current perturbation of the

PTM and is in phase with the positive temperature spikes. With the Hilbert transform it is possible to compute Hilbert Amplitude spectra (HAS) from the IMFs [7]. This is demonstrated in Fig. 3. EMD was applied on magnetics data when a PTM was present (coil T002, JET pulse: #52011).

The obtained spectrum shows only one dominant mode and some signs of intra-wave modulations. A similar behaviour is observed in HAS of ECE data although the noisy data sets aggravate the interpretation. Usually, in Fourier spectra of PTMs up to seven harmonics are observed. These are also visible in wavelet analysis if the popular Morlet wavelet is used since it is Fourier based. A drawback of HAS is that the frequency of the PTM cannot be as easily determined as in FFTs or wavelet scalograms. It is therefore useful to combine several methods to make use of all advantages.




Figure 3: Hilbert amplitude spectrum of coil T002 (JET pulse #52011)

Conclusion

EMD and HAS from single signals thereby consistently supports the assumption that the PTM is indeed a current filament. The magnitude of the current-induced temperature fluctuations can also be deduced and allows an estimate of the influence on the equilibrium.

Acknowledgement

This work was supported by EURATOM and carried out within the framework of the European Fusion Development Agreement. The views and opinions expressed herein do not necessarily reflect those of the European Commission. This work was also supported by grant P19901 of the Austrian Science Funds (FWF).

References

- [1] E. R. Solano, Phys. Rev. Lett. **104**, 185003 (2010)
- [2] H. Koslowski et al., Nuclear Fusion **45**, 201 (2005)
- [3] Ch. Maszl et al., Journal of Nuclear Materials (2011), doi: 10.1016/j.jnucmat.2010.12.245.
- [4] P. Miglicucci et al., Physics of Plasmas **17**, no. 7, 072507, (2010)
- [5] J. R. Myra et al., Physics of Plasmas **14**, no. 10, 102314, (2007)
- [6] N. Commaux et al., Nucl. Fus. **50**, no. 2, 025011, (2010)
- [7] N. E. Huang et al., Proc. R. Soc. Lond. A **454** (1998), 903-995